



Publication number:

0 473 502 A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 91402307.2

© Int. CI.5 A61K 7/48, C11D 3/33

- (22) Date of filing: 23.08.91
- Priority: 24.08.90 JP 223553/90 01.11.90 JP 295678/90
- Date of publication of application: 04.03.92 Bulletin 92/10
- Designated Contracting States:

 DE FR GB IT

- Applicant: SHISEIDO COMPANY LIMITED 7-5-5 Ginza Chuo-ku Tokyo(JP)
- (2) Inventor: Kitamura, Kenji, c/o Shiseido Laboratories 1050, Nippa-cho, Kohoku-ku Yokohama-shi, Kanagawa(JP) Inventor: Nakayama, Yasukazu, c/o Shiseido Laboratories 1050, Nippa-cho, Kohoku-ku Yokohama-shi, Kanagawa(JP) Inventor: Akiyama, Naoe, c/o Shiseido Laboratories 1050, Nippa-cho, Kohoku-ku Yokohama-shi, Kanagawa(JP) Inventor: Kasahara, Eriko, c/o Shiseido Laboratories 1050, Nippa-cho, Kohoku-ku Yokohama-shi, Kanagawa(JP)
- (2) Representative: Rinuy, Santarelli 14, avenue de la Grande Armée F-75017 Paris(FR)
- (A) Washing composition capable of preventing and ameliorating skin irritation.
- (a) A washing composition containing one or more compounds having a protease inhibitory activity and dermatologically acceptable components.

BACKGROUND OF THE INVENTION

1 Field of the invention

10

20

The present invention relates to a novel washing agent. More specifically, it relates to a detergent composition for domestic or business use or a skin cleanser composition comprising one compound or two or more compounds selected from compounds having a protease inhibitory activity, which prevents and ameliorates skin irritation, and at the same time, is harmless to the human body and has a strong washing power.

2. Description of the Related Art

Washing compositions such as cleansers and detergents have been generally employed for removing contaminates or soils attached to the surface or internally of materials and the human body, and may be broadly classified into 1) those used for the human body, 2) those used for clothing and fiber products, 3) and those for washing soft and rigid surfaces.

Among these cleansers, the cleanser for the human body (i.e., skin) is intended to remove the fat and sweat secreted from the skin surface, and other corneum cells detached as a result of metabolism (i.e., dirt), outside contamination attached thereto, and cosmetics.

The surfactant contained in a detergent is a chemical exhibiting a washing power and foaming power which are inherent properties of the detergent, but these are known to have an adverse influence on human skin. A detergent which comes directly into contact with human skin must produce only a weak irritation of the skin or eyes.

For skin cleansers, surfactants with a low irritation effect have been developed to obtain useful products, but there remains a need to improve the washing power and feaming power thereof.

For the washing of the human body, surfactants with a low irritation, effect have been developed to obtain useful products, but there remains a need to improve the washing power and foaming power, and a product which fully satisfies users has not yet been developed. Further, even though the composition has a low irritation effect, it would be more useful if a chemical which treats and ameliorates skin irritation were formulated therein.

On the other hand, currently, for washing objects other than the human body, the surfactants primarily used have a strong washing power but also cause an irritation of the skin or eyes. Although efforts to improve safety have been made by working with the use of protective tools such as gloves, or by way of after-care after completion of the work, the sensitivity of the fingertips may be lost depending the work, and thus the wearing of gloves may be unsuitable in some cases; sometimes skin irritation is reported to be generated by the gloves per se. Also, although the practice of after-care after the completion of work is is important, it is considered more useful if the chemical acts during the work.

Many proteases are known to exist, which maintain the structure and constancy of the skin functions, and the important role thereof thus recently attracted more attention. A protease or proteolytic enzyme is the comprehensive name for enzymes which catalyze a peptide bond by hydrolysis, and such proteases are classified into peptidases and proteinases. The former are enzymes which cleave the peptide bond from the outside of the amino group terminal end or the carboxyl group terminal end of a protein or a peptide chain, and the latter proteinase are enzymes which cleave the specific bond internally of a peptide chain. Such proteinases are known as "protease" in a broad sense, and further, are broadly classified, according to the properties of their active sites, into 1) serine type, 2) thiol (cysteine) type, 3) carboxyl type and 4) metal proteinase, and specific inhibitors exist for the respective enzymes.

SUMMARY OF THE INVENTION

The objects of the present invention are to obviate the above-mentioned problems of the prior art and to provide a washing composition containing one or more compounds having a protease inhibitory activity, which prevents and ameliorates skin irritation, and is harmless to the human body and has a strong washing power.

Other objects and advantages of the present invention will be apparent from the following description. In accordance with the present invention, there is provided a washing composition comprising at least one compound having a protease, inhibitory activity, and dermatologically acceptable components for a conventional washing composition.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present inventors, in view of the state of the art as described above, have made an intensive study of the relationship between various chemicals and skin irritation treatment and amelioration, for treating and ameliorating skin irritation by formulating an effective chemical in the detergent itself, which has not been practiced in the prior art, and consequently, found that a washing composition formulated with a protease inhibitor provides an excellent amelioration of skin irritation, and accomplished the present invention based on thus finding.

The constitution of the present invention is now described as follows.

More specifically, the first aspect of the present invention concerns a detergent for domestic or business use, comprising one or two or more compounds selected from compounds having a protease inhibitory activity.

The detergent composition for domestic or business use of the present invention includes washing cetergents for domestic use, washing detergents for business use (laundry detergents, dry cleaning detergents), washing aids (softeners bleaching agents, glue agents, water repellents), stain remover, carpet shampoo, rust remover, detergents for kitchen (for vegetables, fruits, dish, cocking tools, etc.), oven cleaner, floor polish, floor wax, floor cleaner, floor wax remover, stain remover of floor, window glass detergent, blind cleaner, toilet stool detergent, porcelain detergent, marble detergent, stone wall detergent, wall paper detergent, wall paper and label peeling agent, paint face detergent, paint peeling agent, plastic cleaner, metal face detergent, furniture polisher, slippery agent of door, paper screen, automobile body detergent, car polish, automobile part detergent (type cleaner, inner engine carbon remover, radiator cleaner, methanical part detergent electrical insulator detergent, brake lining cleaner, etc.), aerocraft detergent, aluminum detergent copper and general metal detergent, stainless steel and chromium face detergent, aluminum detergent copper and copper alloy detergent silver detergent, etc.), detergents for press for printing, rolls ink bottle, etc., detergents for dairy industries, tank detergents, discharge pipe, sewage reservoir detergent, leather product detergent, shoe cleaner, shoe polish, spectacle lens cleaner, artificial tooth cleaner, refrigerator detergent, deodorant, road detergent, etc.)

The second aspect of the present invention concerns a skin cleanser composition comprising one or two or more compounds selected from compounds having a protease inhibitory activity.

The skin cleanser composition of the present invention is intended to remove various contaminates attached to the human body, and from animals such as pets and domestic stock.

Specifically, the following compositions may be included in the second aspect of the present invention, but it is not limited thereto.

The compositions include soap, hand cleansers, body shampoos, body lotions, pre-shaving lotions, shaving foams, cleansing creams, cleansing lotions, and pet shampoos.

A protease or proteolytic enzyme is the comprehensive name for enzymes which catalyze a peptide bond by hydrolysis, and such proteases are classified into peptidases and proteinases. The former are enzymes which cleave the peptide bond from the outside of the amino group terminal end or the carboxyl group terminal end of a protein or a peptide chain, and the latter proteinase are enzymes which cleave the specific bond internally of a peptide chain. Such proteinases are known as "protease", in a broad sense, and further, are broadly classified, according to the properties of their active sites, into 1) serine type, 2) thiol (cysteine) type, 3) carboxyl type and 4) metal proteinase, and specific inhibitors exist for the respective enzymes.

The protease inhibitor or the compound having a protease inhibitory activity usable in the present invention includes all the chemical substances capable of inhibiting, reversibly or irreversibly, the hydrolysis activity of the above protease or proteolytic enzyme.

Specifically, the following substances are included.

- (1) Compounds derived from animals or vegetables:
- preferably a bovine pancrea basic trypsin inhibitor, aprotinin, soybean trypsin inhibitor, rimabean professe inhibitor, and corn professe inhibitor.
- (2) Compounds derived from microorganisms:

preferably antipine, plasminostreptin, and further, compounds comprehensively known as leupeptin and represented by the following formula:

$$R_1 - R_2 - R_3 - NH - CH - (CH_2)_3 - NH - CH - NH_2$$

50

55

10

20

 R_1 = CH_3CO , CH_3CH_2CO R_2 = L-Leu, L-IIe, L-Val R_3 = L-Leu, L-IIe, L-Val

(Leu: leucine, lie: scieucine, Val. valine)

(3) Benzamidine and derivatives thereof.

preferably benzamidine, p-aminobenzamidine, m-aminobenzamidine, phenylguanodine, (2R,4R)-4-methyl-1- $[N^2-(3-methyl-1,2,3,4-tetrahydro-8-quinolinesulfonyl)$ -L-arginyl]-2-piperidinecarboxylic acid monohydrate, and dancylarginine N-(3-ethyl-1,5-pentaned yl)amide.

(4) Acetamide and derivatives thereof:

preferably acetamide and 2-phenylacetamide, cyclohexylacetamide.

(5) Guanidine and derivatives thereof:

preferably phenylguanidine and cyclohexylguanidine

(6) ω-amino acids:

Preferably tranexamic acid, p-aminomethylbenzoic acid. 4-aminomethylbicyclo(2.2.2)octane-1-carboxylic acid, 5-{trans-4(aminomethyl)cyclohexyl}-tetrazole, 3-{trans-4(aminomethyl)cyclohexyl-2-ox-opropionate, trans-4-(aminomethyl)cyclohexyl glyoxal monohydrate, and trans-4-(aminomethyl)cyclohexane hydroxamic acid, or substances represented by the following formula wherein the carbon chain has n = 1 - 8:

NH₂(CH₂)_nCOOH

The present invention is not limited to the above, but among these ω -amino acids, particularly good effects can be recognized in ϵ -aminocaproic acid and tranexamic acid, and in p-aminomethylbenzoic acid.

(7) Fluorophosphoric acid and derivatives thereof preferably diisopropylfluorophosphoric acid.

(8) Fluorosulfonic acid and derivatives thereof:

preferably phenylmethanesulfonyl fluoride, and [(p-amidinophenyl)methanesulfonyl fluoride.

(9) Guanidinobenzoic acid and derivatives thereof:

Preferably p-nitrophenyl-p'-guanidinobenzoic acid, 3'.6'-bis(4-guanidinobenzoyloxy)-5-(N'-4-carboxyphenyl)thioure.dospiro[isobenzofuran-1(3H) 9'-(9Hixanthen]-3-one.

(10) Lysine and derivatives thereof:

preferably compounds represented by the following formula:

35

40

5

1 C

15

20

25

30

R₁ = H, Phe-Ala, Ala-Phe

R₂ = OH, CH₂CI

 $R_3 = H$.

45

5*0*

55

(Phe: phenylatanine, Ala: atanine)

The present invention is not limited to the above, but among these lysine and derivatives thereof, a particular good effect is recognized in $R_2 = CH_2CI$

(11) Arginine and derivatives thereof:

preferably the compounds represented by the following formula:

R. = H. D-Phe-Pro. Glu-Gly. Ile-Glu-Gly. Pro-Phe Ala-Phe

 $R_2 = OH, CH_2CI$

 $R_3 = H$

15

5

•0

(Phe: phenyialanine, Pro: proline, Glu:glutamic acid, Gly: glycine, Ile: isoleucine, Ala: alanine). The present invention is not limited to the above, but among these arginind and derivatives thereof, a particularly good effect is recognized in $R_2 = CH_2CI$.

The respective substances as mentioned above have been known only to have protease inhibitory activities, and have not been known to prevent and ameliorate skin irritation.

In the present invention, the compounds having a protease inhibitory activity as mentioned above can be used alone or in any combination thereof.

Also, in the present invention, the amount of the compound having a protease inhibitory activity and formulated in the washing composition is preferably 0.001 to 20% by weight, more preferably 0.001 to 5% by weight. If less than 0.0001% by weight, the effect of the present invention is not obtained, and an amount over 20% by weight is not preferable from the standpoint of preparation and cost.

The "skin irritation treatment and amelioration effect" as mentioned in the present invention means the activity as shown below.

Specifically, this term refers to an amelioration of the disappearance or obscuration of the peeling of the corneum, skin grooving and skin swelling due to the use of the cleaners of the prior art.

In the washing composition of the present invention, in addition to the compounds having a protease inhibitory activity, there can be formulated, if necessary, pharmaceutically or dermatologically acceptable surfactants (anionic, cationic, nonionic, amphoteric, semipolar) aids (builder), abrasiues, fluorescent brightness, bleaching agents, colorants, preservatives, sequestering agents, antistatic agents, precipitation preventives, antioxidants, perfumes, oils, humectants, and chemicals such as antiphlogistic agents, sterilizers, and vitamins within the range which does not impair the effect of the present invention.

As examples of the above-mentioned additives, the anionic surfactants include:

i) fatty acid soap type anionic surfactants represented by the formula:

40 RCOOM

(wherein R represent an alkyl group or alkenyl group having 8 to 18 carbon atoms, M represents one or two or more of alkali metals, organic amines, and basic amino acids);

ii) ether carboxylic acid salt type anionic surfactants represented by the formula:

R(OCH2CH2),OCH2COOM

(wherein R represents an alkyl group or alkylallyl group having 8 to 22 carbon atoms, n is an integer of 1 to 16, and M represents one or two or more of alkali metals, organic amines, and basic amino acids);

iii) N-acylsarcosine salt type anionic surfactants represented by the formula

CH₃ RCONCH₂COOM

55

45

50

(wherein R represent an alkyl group or alkenyl group having 8 to 18 carbon atoms, and M represents one or two or more of alkali metals, organic amines, and basic amino acids);

v) anionic surfactants having a

-COOT group

10

30

35

in the formula represented by condensed products of higher fatty acids and amino acids such as N-acylglutamic acid salts represented by the formula:

CH2CH2COOM RCONHCHCOOM

(wherein R represents an alkyl group or alkenyl group having 8 to 18 carbon atoms, and M represents one or two or more of alkali metals, organic amines, and basic amine acids).

More specifically, anionic surfactants can include base materials for soap; fatty acid scaps such as sodium laurate, sodium palmitate or the like; higher alkyl sulfate salts such as sodium lauryl sulfate, potassium lauryl sulfate or the like; alkyl ether sulfate salts such as triethanolamine polyoxyethylene (hereinafter abbreviated as POE) laury! sulfate, sodium POE laury! sulfate or the like; N-acy!sarcosine such as sodium lauroylsarcosinate or the like; higher fatty acid amide sulfonic acid salts such as sodium Nmyristoyl-N-methyltaurine, sodium coconut oil fatty acid methyltauride, sodium lauryl methyltauride or the like; phosphate salts such as sodium POE oleyi ether phosphate, and POE stearyl ether phosphoric acid or the like; alkythydroxycarboxylic acid salts such as sodium laurylhydroxy ether carboxylate or the like; 20 sulfosuccinic acid salts such as sodium di-2-ethylhexylsulfosuccinate, sodium monolauroyl moncethanolamide polyoxyethylene sulfosuccinate, and sodium lauryl polypropylene glycol sulfosuccinate or the like; alkylbenzene sulfonic acid salts such as sodium linear dodecylbenzene sulfonate, triethanolamine linear dodecylbenzene sulfonate, and linear dodecylbenzene sulfonic acid or the like; N-acylglutamate salts such as monosodium N-lauroylglutamate, disodium N-stearoylglutamate, and monosodium N-myristoyl-L-25 glutamate or the like; higher fatty acid ester sulfuric acid salts such as sodium hardened coconut oil fatty acid glycerine sulfate or the like; sulfated oil such as Turkey red oil; POE alkyl ether carboxylic acids, POE alkyfallyl ether carboxylic acid salts, a-olefinsulfonic acid salts, higher fatty acid ester sulfonic acid salts. secondary alcohol sulfonate salts, higher fatty acid alkylolamide sulfate salts, sodium lauroyl moncethanolamide succinate, di-triethanolamine N-paimitoylaspartate, and sodium caseinate, and so on.

The cationic surfactants include monoalkyl type quaternary ammoniums salts represented by the formula:

$$\begin{bmatrix} R_2 \\ R_1 - N - R_4 \\ R_3 \end{bmatrix} + X^-$$

(wherein R- represents an alkyl group or alkenyl group having 12 to 22 carbon atoms, R_2 , R_3 and R_4 represent methyl or ethyl group, X represents a halogen atom or a methyl sulfate residue), and further, aiiphatic amine salts, aromatic quaternary ammonium salts, pyridinium salts, and imidazolinium salts.

The nonionic surfactants can include glycerine fatty acid esters, sorbitane fatty acid esters sorbitol fatty acid esters, sucrose fatty acid esters, polyoxyethylene (hereinafter called POE) sorbitane fatty acid ester, polyoxyethylene glyccl fatty acid esters, POE alkyl ethers, POE alkylphenyl ethers, POE hardened castor oil derivatives, mannitol hydroxyfatty acid ethers, and alkylglycoside fatty ethers

The amphoteric surfactants can include carboxybetaines such as N,N-dimethyllauryl-N-carboxymethylammonium betaine, N N-dimethyl-N-oleyl-N-carboxymethylammonium betaine, and lauryldimethyl-aminoacetic acid or the like imidazoline derivatives such as 2-lauryl-N-carboxyethyl-N-hydroxyethylimidazolinium betaine, 2-lauryl-N-carboxymethyl-N hydroxyethylimidazolinium betaine, 2-undecyl-N,N,N-(hydroxyethylcarboxymethyl)-2-imidazoline sodium, and 2-cocoyl-2-imidazolinium hydroxide-1-carboxyethyloxy-2-sodium or the like aminocarboxylic acid salts such as sodium N-coconutalkyl- β -aminopropionate and sodium N-coconutalkyl- β -iminodipropionate or the like, and sulfobetaine and aminobetaine

The semi-polar surfactants can include lauryldimethylamine oxide, stearyldimethylamine oxide, and bis-(2-hydroxyethyl)lauryl-amine oxide.

As the propellant, all propellants which can be used in aerosol products in general are applicable. Specifically, there include fluorinated hydrocarbons such as Freon 11 (registered trade mark) Freon 12

(registered trade mark). Freon 21 (registered trade mark), Freon 113 (registered trade mark), and Freon 114 (registered trade mark), inquefied petroleum gas (L. P. G.) which is a mixture of propane, isobutane, nebutane, and a compressed gas such as carbon dioxide gas and nitrogen gas. These gases can be used either alone or as a mixture of two or more kinds thereof, and the amount formulated is generally 2 to 20% by weight. With an amount of 2% or less, the internal pressure may be lower, and thus the stock liquid may not be properly emitted upon use, and on the other hand, further improved results cannot be obtained if 20% or more of the gas is employed; conversely, the internal pressure may become too high.

Any conventionally used water-insoluble oil may be employed, for example, higher alcohols such as cetyl alcohol, stearyl alcohol, and cholesterol or the like; higher fatty acids having C₈ - C₂₂ carbon atoms such as lauric acid, myristic acid, palmitic acid, stearic acid, oleic acid, and isostearic acid or the like, waxes such as solid paraffin, microcrystalline wax, polyethylene wax, canderilla wax, beeswax, hardened castor oil, carunauba was, and barico wax or the like; animals and vegetable oils such as tallow, lard, sheep oil, squalane, coconut oil, palm oil, palmkernel oil, soybean oil, olive oil, cottonseed oil, jojoba oil, castor oil, and lanolin, mineral oils such as fluid paraffin and petrolatum; and synthetic oils such as trimethylpropane triisostearate, isopropyl myristate, giycerol tri-2-ethylhexanate, pentaerythritol tetra-2-ethylhexanate, silicone oil, and polyoxyethylene polyoxypropylene pentaerythritol ether.

Any conventionally used polyhydric alcohol may be employed, for example, propylene glycol, dipropylene glycol, glycerine, 1.3-butylene glycol, polyethylene glycol, polyoxyethylene methylglycoside ether, polyoxyalkylene diglyceryl ether, polyoxyalkylene decaglyceryl ether, polyoxyalkylene pentaerythritol ether, sorbitol, maltitol, lactose, and D-mannitol.

Further, there can be included sterilizers such as cetylpyridinium chloride, benzetonium chloride, decalinium chloride, benzalkonium chloride, chlorohexydine gluconate, carbanilide, phenol, and halogenated salicylanilide, alkalis such as caustic potash and ammonia, lower alcohol such as ethanol, and humectants such as mucopolysaccharides and pyrrolidone carboxylic acid salts, etc. Other humectants are propylene glycol, dipropylene glycol, glycerine, 1,3-butylene glycol, polyethylene glycol, polyoxyethylene methylglycoside ether, polyoxyalkylene diglyceryl ether, polyoxyalkylene pentaerythritol ether, sorbitol, maltitol, lactose, D-mannitol, mucopolysaccharide, and pyrrolidone carboxylic acid salts.

The abrasives usable in the detergent composition for domestic or business use include silicates such as sodium metasilicate or the like, salts such as sodium sulfate and sodium carbonate, silica, borax, talc, diatomaceous earth, bentonite, colloidal clay, fluorite, quartz, and sand.

The washing composition according to the present invention may be in any desired form, as long as it is in the form suitable for accomplishing the object of providing a detergent, as exemplified by a liquid (lotion, milky lotion), cream, solid, fine granules, and powder.

The washing composition of the present invention can be used while washing, and prevents and ameliorates skin irritation, and at the same time, is harmless to the human body and has a strong washing power.

EXAMPLES

40

The present invention will now be further illustrated by, but is by no means limited to, the following Examples and Actual Use Examples.

Prior to the Examples, the test methods and evaluation methods practiced for clarifying the prevention of skin irritation and amelioration of skin irritation effects of the compound having a protease inhibitory activity according to the present invention are explained.

Actual Use Test I-1

Sixty housewives 40 to 60 years old and susceptible to "skin irritation" when continuously using conventional synthetic detergents in the kitchen were selected as the subjects to be tested and divided into 6 groups each having 10 members. For the subjects to be tested of 5 groups, the respective Examples shown in Table I-1 were used, and for the subjects to be tested of the remaining one group, Comparative Example I-1 was used. Continuous use tests were conducted for 4 weeks, and thereafter, the state of the skin of the back of the hands was observed by the replical method, and evaluated according to the judgement standards shown in Table I-2. Each recipe was prepared according to conventional methods, to obtain a synthetic detergent for kitchen.

35 40 45	30	25	20	15	10	5
		Table I-1				
			Example			Comparative Example
	1-1	1-2	I-3	7−I	1-5	I-1
Sodium POE (3 mole) alkyl ether sulfate	15.0	15.0	15.0	15.0	15.0	15.0
POE (15 mole) alkyl ether	5.0	5.0	5.0	5.0	5.0	5.0
Coconut fatty acid ethanol amide	6.0	0.9	6.0	0.9	6.0	6.0
Ethanol	1.0	1.0	1.0	1.0	1.0	1.0
Perfume	. s. p	8. b	.s.p	. s . p	q.s.	d.s.
Dye	q.s.	. s . b	q.s.	.s.	q.s.	q.s.
Purified water	balance	balance	balance	balance	balance	balance
Tranexamic acid	1.0	1	ı	t	1	•
Leupeptin	1	0.5	ı	ı	ı	ŧ
Tosylarginine	ı	ı	3.0	1	1	ı
Tosyllysyl chlormethyl ketone	ı	ı	1	0.05	ł	•
Soybean trypsin inhibitor	ı	ŧ	1	1	0.1	,

Table I-2

5	Score	Evaluation	Remarks
	1	Disappearance of skin groove, skin rise Peeling of wide range of corneum	Irritated skin
10	2	Obscure skin groove, skin rise Partial peeling of corneum	Î
	3	Skin groove, skin rise recognized, but flat	
	4	Clear skin groove, skin rise	
15	5	Skin rise, skin groove clear and regular	Beautiful skin

Table I-3

			Comparative Example						
Replica score	I-1	I-2	I-3	I-4	1-5	I-1			
1	0	0	. 0	0	0	2			
2	o	0	0	0	0	2			
3	0	0	1	2	1	5			
4	2	1	1	1	4	1			
5	8	9	8	7	5	0			

The numerals in the Table indicate the number of the persons tested who exhibited the respective evaluation scores.

As shown by the results in Table I-3, the detergent compositions of the present invention (Examples I-1 - I-5) exhibited a superior effective to that of the detergent composition of the Control (Comparative Example I-1).

Example I-6 Liquid detergent for clothing

50

45

20

25

30

35

	(1)	Na POE (3 mole) lauryl ether sulfate	15.0
	(2)	POE (15 mole) alkyl ether	20.0
£	(3)	Distearyl dimethylammonium chloride	2.0
	(4)	Aprotinin	0.5
	(5)	Bleaching agent	q.s.
	(6)	Purified water	balance
10	Exar	nple I-7 Powder detergent for clothing	ſ
	(1)	LAS-Na	15.0
	(2)	Na ₂ SO ₄	30.0
	(3)	2-Phenylacetamide	2.0
15	(4)	CMC (66%)	1.5
	(5)	Sodium metasilicate (anhydrous)	20.0
	(5)	Fluorescent brightener	0.2
20	(7)	Na ₂ CO ₃	balance
	<u>Exam</u>	nple I-8 Solid soap	
	(1)	Tallow	20.0
	(2)	Coconut oil	12.0
25	(3)	Castor oil	5.0
	(4)	Olive oil	3.0
	(5)	Caustic potash	6.0
	(6)	Ethanol	20.0
30	(7)	Glycerine	5.0
	(8)	Sucrose	10.0
	(9)	Cyclohexylguanidine	3.0
	(10)	EDTA	0.1
35	(11)	Perfume	q.s.
	(12)	Dye	q.s.
	(13)	Purified water	balance
40	Exam	ple I-9 Liquid soap	
-	(1)	Lauric acid	3.0
	(2)	Myristic acid	7.0
	(3)	Palmitic acid	3.0
45	(4)	Oleic acid	2.5
	(5)	Lauroyl diethanolamide	6.0
	(6)	Propylene glycol	11.0
	(7)	Glycerine	4.0
50	(8)	Sucrose	5.0

(9)	Caustic potash	3.0
(10)	EDTA	0.1
(11)	p-Aminobenzimidine	1.0
(12)	Perfume	q.s.
(13)	Purified water	balance

The respective compounds used in the following Examples were the compounds represented by the formula shown below, in which R_1 , R_2 , R_3 are respectively those shown in Table I-4.

20

15

5

Table I-4

R ₁ R ₂	R ₃
Compound I-1 D-Phe-Pro CH ₂ C1	so ₂ -CH ₃
Compound I-2 Glu-Gly CH ₂ Cl	so ₂ CH ₃
(Phe: phenylalanine, Pro: proli acid, Gly: glycine) <u>Example I-10</u> Detergent for kitch	
(1) Na α-Oleylsulfonate	20.0
(2) POE (15 mole) alkyl ether	8.0
(3) Laurylamide propylbetaine	5.0
(4) Ethanol	1.5
(5) Compound I-1	3.5
(6) Dye	q.s.
(7) Perfume	q.s.
(8) Purified water	balance
Example I-11 Detergent for kitch	en
(1) Monosodium N-lauroylglutamat	e 25.0
(2) POE (15 mole) alkyl ether	5.0

	(3)	Lauryldimethylamine oxide	8.0
	(4)	Ethanol	1.0
5	(5)	Compound I-2	7.0
	(6)	Dye	q.s.
	(7)	Perfume	q.s.
16	(8)	Purified water	balance

All of the detergent compositions of Examples I-6 - I-11 were found to provide a superior skin irritation and skin irritation amelioration effect, to be harmless to the human body, and to have a strong washing power.

Actual Use Test II-1

Sixty men 30 to 50 years old and susceptible to "razor irritation" were selected as the subjects to be tested and divided into 6 groups each having 10 members. For the subjects to be tested of 5 groups, the respective Examples shown in Table II-1 were used, and for the subjects to be tested of the remaining one group. Comparative Example II-1 was used. Continuous use tests were conducted for 4 weeks, and thereafter, the state of the skin back of the hand was observed and evaluated according to the standards given in Table II-2. Each recipe was prepared according to conventional methods, to obtain an aerosol shaving foam.

10		Example Example	II-3 II-4 II-5 II-1	4.2 4.2 4.2	3.5 3.5 3.5	8.0 8.0 8.0	3.0 3.0 3.0	q.s. q.s. q.s. q.s.	q.s. q.s. q.s. q.s.	balance balance balance	0.9 6.0 6.0				- 5.5 -	
25	Table II-1		11-2	4.2	3.5	8.0	3.0	d.s.	• 6 • 	balance	6.0	•	1.2	ı	ı	
30	₽I		11-1	e 4.2	3.5	8.0	3.0	q.s.	9.5	balance	6.0	0.3	ı	•	1	
35				C ₁₈ alkyltrimethylammonium chloride	lutamate	cetic acid								yl ketone		7
40				yltrimethylam	Potassium myristoylglutamate	Lauryldimethylaminoacetic acid betaine	ne			d water	ant (L.P.G.)	in	mic acid	Tosyllysyl chlormethyl ketone	ginine	Sovhean trynsin inhihitor
45				Cl8 alky	Potassi	Lauryld: betaine	Glycerine	Perfume	Dye	Purified wat	Propellant (Leupeptin	Tranexamic	Tosylly	Tosylarginin	Sovbean

(The numerals in the Table show I by weight.)

50

Table II-2 Amelioration effect on razor irritation

Evaluation	Judgement standards							
Remarkably effective	Razor	irritation	disappeared					
Effective	•	•	greatly ameliorated					
Slightly effective	•		slightly ameliorated					
Ineffective	•	•	not changed					
Worsened	•	•	worsened					

Table II-3

Judgement results by		Comparative Example				
visual observation	II-1	11-2	II-3	11-4	11-5	II-1
Remarkably effective	8	9	6	8	6	0
Effective	2	ĭ	3	1	2	1
Slightly effective	0	0	1	1	1	2
Ineffective	0	0	0	0	1	7
Worsened	0	0	0	0	0	0

The numerals in the Table show the number of the persons indicating the respective judgment results. As shown by the results in Table II-3, the skin cleanser compositions of the present invention (Examples II-1 - II-5) exhibited a superior effective to that of the skin cleanser composition of the Control (Comparative Example II-1).

45 Actual Use Test II-2

Thirty women 25 to 40 years old and susceptible to skin irritation were selected as the subjects to be tested and divided into three groups each having 10 members. For the subjects to be tested of two groups, the respective Examples shown in Table II-4 were used, and for the remaining one group, the Comparative Example II-2 was used, and face washing was practiced. The tests were practiced by continuous use for 4 weeks, and after completion of the test, the skin state of the cheek portion of the face was judged by the replical method. The judgment was conducted according to the standards shown in Table II-5, and the results were represented as the average of the panel of 10 members. The manner in which contaminates such as cosmetics, etc., were removed was judged by a self-evaluation by the panel members, following the standards shown in Table II-6. Each recipe was prepared according to conventional methods, to obtain a cleansing foam.

Table II-4

	Exa	mple	Comparative Example
	II-6	II-7	II-2
Lauric acid	8.0	8.0	8.0
Myristic acid	12.0	12.0	12.0
Palmitic acid	4.0	4.0	4.0
Stearic acid	4.0	4.0	4.0
Glycerine monostearate	1.5	1.5	1.5
POE glycerylmonostearate (60E.O.)	2.0	2.0	2.0
Glycerine	2.5	2.5	2.5
Polyoxyethylene glycol (M.W. 500)	8.0	8.0	8.0
Sorbitol solution	5.0	5.0	5.0
Caustic potash	5.5	5.5	5.5
Purified water	balance	balance	balance
p-Aminobenzamidine	0.5	-	-
Tosyllysine	-	2.0	-

C

Table II-5

Scor	Evaluation	Remarks
1	Disappearance of skin groove, skin rise Peeling of wide range of corneum	Irritated skin
2	Obscure skin groove, skin rise Partial peeling of corneum	
3	Skin groove, skin rise recognized, but flat	
4	Clear skin groove, skin rise	Į
5	Skin rise, skin groove clear and regular	Beautiful skin

Table II-6

25	Evaluation		Conten	ts	
	©	80% or more of removal of cont		satisfied with	good
30	0	607	•	•	
	Δ	407	•	•	
35	x	Less than 40%	•	•	

Table II-7

	Exam	ple	Comparative Example
	II-6	II-7	I I- 2
Replica score	4.8	4.7	2.9
Extent of contaminate removal	o	©	©

As shown by the results given in Table II-7, the skin cleanser compositions (Examples II-6 and II-7) exhibited a superior effective to that of the skin cleanser composition of the Control (Comparative Example 55 II-2).

Example II-8 Solid soap

		*	by weight
5	(1)	Tallow	20.0
3	(2)	Coconut oil	12.0
	(3)	Castor oil	5.0
	(4)	Olive oil	3.0
10	(5)	Caustic potash	6.0
	(6)	Ethanol	20.0
	(7)	Glycerine	5.0
15	(8)	Sucrose	10.0
	(9)	Cyclohexylguanidine	3.0
	(10)	EDTA	0.1
20	(11)	Perfume	q.s.
	(12)	Dye	q.s.
	(13)	Purified water	balance
	Exam	ple II-9 Liquid soap	
25	(1)	Lauric acid	3.0
	(2)	Myristic acid	7.0
	(3)	Palmitic acid	3.0
30	(4)	Oleic acid	2.5
	(5)	Lauroyl diethanolamide	6.0
	(6)	Propylene glycol	11.0
35	(7)	Glycerine	4.0
	(8)	Sucrose	5.0
	(9)	Caustic potash	3.0
	(10)	EDTA	0.1
10	(11)	p-Nitrophenyl-p'-guanidinobenzoic ac	id 0.1
	(12)	Perfume	q.s.
	(13)	Purified water	balance

The respective compounds used in the following Examples were the compounds represented by the formula shown below, in which R_1 , R_2 , R_3 are respectively those shown in Table II-4.

55

Table II-8

:		R ₁	R ₂	R	3
	Compound II-1	D-Phe-Pro	CH ₂ C1	so ₂ \sqrt{C}	CH ₃
70	Compound II-2	Glu-Gly	СН ₂ С1	so ₂ ~	Сн3
	Compound II-3	Ile-Glu-Gly	CH ₂ Cl	so ₂ -	CH ₃
15	Compound II-4	Pro-Phe	CH ₂ C1	so ₂	CH3
26	(Phe: phenyla acid, Gly: c Example II-10	glycine, Ile	: isole		glutamic
				%]	by weight
25	(1) Solid pa				3.5
	· · ·	stalline wa	x		8.0
	(3) Beeswax				5.0
30	(4) Petropla				1.0
	(5) Fluid pa				43.0
	· · · -	monooleate			4.0
	•	mole) monoo	leat e		0.5 5.0
35	(8) Compound (9) Perfume	15 11-1			
	(10) Purified	l water			q.s. balance
	Example II-11		d tyme c'	leansing	
40	Example 11-11	_ Linui Silie	a cype c.	-	by weight
	(1) Stearic	acid		0 1	3.0
	(2) Cetyl al				1.0
	(3) Petrolat				4.5
45	(-, 1001010	· -			

50

	(4)	Fluid paraffin	11.0
	(5)	POE (20 mole) oleyl alcohol ether	1.5
5	(6)	POE (5 mole) sorbitane monolauric	1.0
		acid ether	
	(7)	Triethanolamine	1.0
10	(8)	Propylene glycol	6.0
, 0	(9)	Compound II-2	1.0
	(10)	Perfume	q.s.
	(11)	Purified water	balance
15	Exam	nple II-12 Non-emulsified type cleans:	ing lotion
			y weight
	(1)		10.0
20	(2)	Dipropylene glycol	6.0
	(3)	N, N-dimethyl-lauryl-N-carboxymethyl-	2.0
		ammoniumbetaine	
25	(4)	POE (20 mole) sorbitane monolauric	3.0
13		acid ether	
	(5)	Ethyl alcohol	10.0
	(6)	Compound II-3	5.0
30	(7)	Perfume	q.s.
	(8)	Purified water	balance
	<u>Exam</u>	ple II-13 Aerosol shaving foam	
35	(Fil	ling recipe) % h	y weight
	(1)	Stearic acid	5.5
	(2)	Purified coconut fatty acids	1.5
40	(3)	Glyceryl monostearate	5.0
	(4)	Glycerine	10.0
	(5)	Triethanolamine	4.0
	(6)	Compound II-4	0.05
45	(7)	Perfume	q.s.
	(8)	Purified water	balance
	(Fil:	ling recipe)	
50	(1)	Stock solution	94.0
	(2)	LPG	6.0

Examples II-8 to II-13 were found to perfectly harmless and to have a strong washing power.

Claims

1. A washing composition comprising at least one compound having a protease inhibitory activity and

dermatologically acceptable components for a conventional washing composition

- A washing composition as claimed in claim 1, wherein said compound is contained in an amount of 0.0001% to 20% by weight cased on the total amount of the composition.
- 3. A washing composition as claimed in claim 1 or 2 wherein said compound is at least one compound derived from animals or vegetables selected from the group consisting of a bovine pancrea basic trypsin inhibitor, aprotinin, soybean trypsin inhibitor, tima bean protease inhibitor, and corn protease inhibitor.
- 4. A washing composition as claimed in claim 1 or 2 wherein said compound is at least a derived from microorganisms component selected from the group consisting of antipine, plasminostreptin and compounds comprehensively known as leupeptine represented by the following formula:

$$R_1 - R_2 - R_3 - NH - CH - (CH_2)_3 - NH - CH - NH_2$$

CHO + NH₂

- R· = CH₃CO, CH₃CH₂CO
- R₂ = L-Leu, L-lle, L-Val
- R₃ = L-Leu, L-IIe, L-Val

(Leu: leucine, Ile: isoleucine, Val: valine)

- 25 5. A washing composition as claimed in claim 1 or 2 wherein said compound is at least one compound selected from the group consisting of benzamidine, p-aminobenzamidine, m-aminobenzamidine, phenylguanidine, (2R,4R)-4-methyl-1-[N²-(3-methyl-1,2,3,4-tetrahydro-8-quinolinesulfonyl)-L-arginyl]-2-piperidinecarboxylic acid monohydrate, and dancylarginine N-(3-ethyl-1,5-pentanediyl)amide.
- 30 6. A washing composition as claimed in claim 1 or 2 wherein said compound is at least one compound selected from the group consisting of acetamide and the derivatives thereof.
 - 7. A washing composition as claimed in claim 1 or 2 wherein said compound is at least one compound selected from the group consisting of guanidine and the derivatives thereof.
 - 8. A washing composition as claimed in claim 1 or 2 wherein said compound is at least one ω-amino acid selected from the group consisting of tranexamic acid, p-aminomethylbenzoic acid, 4-aminomethylbicyclo(2,2,2)-octane-1-carboxylic acid 5-[trans-4(aminomethyl)-cyclohexyl]tetrazole, 3-[trans-4(aminomethyl)cyclohexyl-2-oxopropionate, trans-4(aminomethyl)cyclohexyl glyoxal monohydrate, and trans-4(aminomethyl)cyclohexane hydroxamic acid, or substances represented by the following formula wherein the carbon chain has n ≈ 1 8):

NH₂(CH₂)_nCOOH.

- 9. A washing composition as claimed in claim 1 or 2 wherein said compound is at least one component selected from the group consisting of fluorophosphoric acid, fluoro sulfonic acid, and the derivatives thereof.
- 10. A washing composition as claimed in claim 1 or 2 wherein said compound is at least one component selected from the group consisting of p-nitrophenyl-p'guanidinobenzoic acid, 3',6'-bis(4-guanidinobenzoyloxy)-5-(N'-4-carboxyphenyl)thioureidespiro[isobenzofuran-1(3H) 9'-(9H)xanthen]-3-one.
- 55 11. A washing composition as claimed in claim 1 or 2 wherein said compound is at least one component selected from the group consisting of compounds represented by the following formula:

15

20

35

40

45

$$R_1$$
-NH-(CH₂)₄-CH-CO-R₂
NH
R₃

R. = H. Pho-Ala, Ala-Phe

R₂ = OH, CH₂CI

 $R_3 = H$

5

10

15

25

30

35

so₂ - CH₃

(Pre: phenylalanine, Ala: alanine)

12. A washing composition as claimed in claim 1 or 2 wherein said compound is at least one component selected from the group consisting of the compounds represented by the following formula:

R. = H. D-Phe-Pro. Glu-Gly. Ile-Glu-Gly, Pro-Phe. Ala-Phe

R₂ = OH, CH₂CI

 $R_3 = H$.

$$so_2 \leftarrow \bigcirc \rightarrow cH_3$$

(Phe: phenylalanine, Pro: proline, Glu: glutamic acid, Gly: glycine, Ile: isoleucine, Ala: alanine)

45

40

50

EP 91 40 2307

		SIDERED TO BE RELE		
Category		vith indication, where appropriate. Ilevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CI.5)
×	FR-A-2 216 986 (BAYER	AG)	1-3	A 61 K 7 48
	* page 2, line 3 - line 15 * *			C 11 D 3 33
Α		5, line 1 - line 6; claims 1,2;	4	
	example 4)			
×	DE-A-1 939 419 (BAYER	AG)	1-3	
	the whole document.		13	
		- - -		
Х	FR-A-2 016 080 (HOECH	ST AG)	1,2.8	
Α	FR-A-2 016 080 ()	-		
- î	* claims; examples **		11	
	-,, -, -, -, -, -, -, -, -, -, -, -,			
X	DE-A-1 944 568 (HENKE	•	1.2.6	
	* page 4, line 6 - line 18; cl	aim 1; examples 3,4 **		
×	US-A-4 566 985 (L.C. BR	HNO ET ALS	105	
	*claim 1; examples **	ONO ET AL.)	1.2,5	
			ļ	
×	DE-A-2 401 752 (UNILEV	ER)	1,2,8	TECHNICAL FIELDS
İ	* claims 1,2 * *			SEARCHED (Int. CI.5)
×	WORLD PATENTS INDEX	 – – LATEST Derwent Publications 	1,2,8	C 11 D
· ·	Ltd., London, GB; AN 85-22		1,2,0	A 61 K
į	& JP-A-60 146 815 (LION (
	*abstract **			
x	WORLD DATENTS INDEX	 LATEST Derwent Publications 		
	Ltd., London, GB; AN 81-03		1.2.9	
ł	& JP-A-55 145 798 (SANY)	O CHEM.) 14 November 1980		
	*abstract **]	
1				
1				
1				
İ				
	The present search report has t	peen drawn up for all claims	-	
	Place of search	Date of completion of search	' 	Examiner
	The Hague	28 November 91		_
	CATEGORY OF CITED DOCU			GRITTERN A.G.
	articularly relevant if taken alone	the t	iling date	ent, but published on, or after
de	articularly relevant if combined with ocument of the same catagory		iment cited in the iment cited for ot	
0: no	chnological background on-written disclosure			patent family, corresponding
P: in	termediate document	doci	iment	e remmy, corresponding



EUROPEAN SEARCH REPORT

Application Number

EP 91 40 2307

D	OCUMENTS CONS	IDERED TO BE RELEV	ANT	Γ	
Category		ith indication, where appropriate, evant passages		elevant i claim	CLASSIFICATION OF THE APPLICATION (Int. CI.5)
A	FR-A-2 457 891 (AJINOM examples 30,33 **		+	1.12	TECHNICAL FIELDS SEARCHED (Int. Ci.5)
	The present search report has	•			
	Place of search	Date of completion of search	}		Examiner
Y : A : O : P ·	The Hague CATEGORY OF CITED DOCI particularly relevant if taken alone particularly relevant if combined will document of the same catagory technological background non-written disclosure intermediate document theory or principle underlying the in-	th another D: doct L: doct 8 . men	filing di ument d ument d	ate cited in the cited for o	GRITTERN A.G. ent, but published on, or after e application ther reasons patent family, corresponding

•			
			_